

Featured Correspondence

Effects of glycoprotein IIb/IIIa blockers

To the Editor: The meta-analysis by Hernandez *et al*¹ showed that the reduction of death or non-fatal myocardial infarction with IIb/IIIa inhibitors in patients with an acute coronary syndrome (ACS) was independent of patient age. The trials included were performed before widespread administration of clopidogrel in ACS.

Clopidogrel has been shown to be beneficial in ACS; its administration is a class I recommendation.² Glycoprotein IIb/IIIa inhibition is a class I recommendation only in patients with planned percutaneous coronary intervention (PCI) in the absence of clopidogrel, but is a class IIa recommendation in patients already treated with clopidogrel.²

An analysis of the National Registry of Myocardial Infarction-4 suggested that routinely giving a glycoprotein IIb/IIIa inhibitor to patients with a non-ST-elevation ACS treated with clopidogrel might not be justified, especially if PCI were not performed.³ This may be particularly true in elderly patients, since virtually all studies indicate that elderly patients are more likely to bleed than younger patients; bleeding has been shown to be an independent predictor of mortality in both ACS and PCI patients.⁴

An analysis of the ISAR-REACT 2 trial showed that abciximab did not reduce the 30-day incidence of death or any other component of major adverse cardiac events in older patients with a non-ST-elevation ACS treated with clopidogrel; in fact, it seemed to be harmful.⁵ We are currently analysing the relationship between age and outcome in three other ISAR studies.

In conclusion, while we applaud the excellent analysis of the authors, we caution that the results of their meta-analysis may not be generalisable to all patients with ACS, particularly the elderly pretreated with clopidogrel.

Arun Kalyanasundaram, Peter Berger
Geisinger Medical Center, Danville, PA, USA

Adnan Kastrati
German Heart Centre, Munich, Germany

Correspondence to: Dr A Kalyanasundaram, 21-60, Department of Cardiology, 100 North Academy Drive, Danville, PA 17822, USA; aksundaram1@geisinger.edu

Competing interests: None declared.

References

- Hernández AV, Westerhout CM, Steyerberg EW, *et al*. Effects of platelet glycoprotein IIb/IIIa receptor blockers in non-ST segment elevation acute coronary syndromes: benefit and harm in different age subgroups. *Heart* 2007;93:450-5.
- Braunwald E, Antman EM, Beasley JW, *et al*. and American College of Cardiology. American Heart Association, Committee on the Management of Patients With Unstable Angina. ACC/AHA 2002

guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction: summary article, a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients With Unstable Angina). *Circulation* 2002;106:1893-900.

- Bromberg-Marín G, Marín-Neto JA, Parsons LS, *et al*. National Registry of Myocardial Infarction-4. Effectiveness and safety of glycoprotein IIb/IIIa inhibitors and clopidogrel alone and in combination in non-ST-segment elevation myocardial infarction (from the National Registry of Myocardial Infarction-4). *Am J Cardiol* 2006;98:1125-31.
- Kinnaird TD, Stabile E, Mintz GS, *et al*. Incidence, predictors, and prognostic implications of bleeding and blood transfusion following percutaneous coronary interventions. *Am J Cardiol* 2003;92:930-5.
- Ndrepepa G, Kastrati A, Mehilli J, *et al*. Age-dependent effect of abciximab in patients with acute coronary syndromes treated with percutaneous coronary intervention. *Circulation* 2006;114:2040-6.

The authors' reply: We thank Drs Kalyanasundaram, Kastrati and Berger for their interesting comments about our article.

Patients included in our subgroup meta-analysis were not routinely scheduled for an early coronary revascularisation, so our results are not directly comparable with the ISAR trials. Patients did not receive clopidogrel as background medication at the time of admission, but almost all received dual antiplatelet treatment at the time of percutaneous coronary intervention (PCI). We agree that clopidogrel is recommended in patients with non-ST segment elevation ACS (NSTEMI) whether or not an invasive approach is planned.¹ The concern, however, is the timing of clopidogrel administration, which continues to be debated, with many cardiologists electing to withhold clopidogrel until angiography because of concerns about bleeding with bypass surgery.² This approach is consistent with the treatment strategy applied in the trials included in our meta-analysis.

We agree that troponin-negative patients with moderate-to-high risk NSTEMI seem to have less benefit than troponin-positive patients, as we have shown in a previous post hoc subgroup analysis from our dataset.³ We acknowledge, however, that the analysis was a retrospective subgroup analysis with poor power.

We appreciate the major contribution that the ISAR trials have made to clinical practice, but they differ significantly from trials included in our meta-analysis. It is critical to avoid direct extrapolation from trials testing adjunctive treatments for PCI (as in the ISAR trials) to trials evaluating strategies of initial medical treatment with PCI performed if needed some time later (in many cases several days later). It is also particularly important to avoid scientifically inappropriate subgroup analyses, in which the subgroup is formed at variable times after the treatment allocation.⁴

In the patchwork of evidence from clinical trials several strategies have emerged as viable. Our meta-analysis was meant to provide a context for clinicians choosing a strategy of initial medical treatment followed by consideration of angiography at a later date with deferred initiation of clopidogrel. We agree that our results should not be directly extrapolated beyond that point. In general, our analysis in the severe subset of our patients with NSTEMI shows a similar trend to the age analysis of the ISAR-REACT 2 trial. We look forward to seeing the results of the analysis of the relation between age and outcomes in the four ISAR trials. Furthermore, the continuing EARLY-ACS trial has enrolled over 6000 patients in a trial directly testing the role of eptifibatide in the setting of "upstream" clopidogrel use, with an emphasis on enrolment of an elderly population.⁵

We welcome further interchange and appropriate contrasts of clinical trials carried out in different circumstances in different populations.

Adrian V Hernandez, Cynthia M Westerhout, Ewout W Steyerberg, John P A Ioannidis, Hector Bueno, Harvey White, Pierre Theroux, David J Moliterno, Paul W Armstrong, Robert M Califf, Lars C Wallentin, Maarten L Simoons, Eric Boersma
Department of Quantitative Health Sciences/Wb4, Cleveland Clinic, Ohio, USA

Correspondence to: Dr A V Hernandez, Department of Quantitative Health Sciences/Wb4, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195, USA; hernana1@ccf.org

Competing interests: None declared.

References

- Bassand JP, Hamm CW, Ardissino D, *et al*. Guidelines for the diagnosis and treatment of non-ST-segment elevation acute coronary syndromes. The Task Force for the Diagnosis and Treatment of Non-ST-Segment Elevation Acute Coronary Syndromes of the European Society of Cardiology. *Eur Heart J* 2007;28:1598-60.
- Mehta RH, Roe MT, Milgund J, *et al*. Acute clopidogrel use and outcomes in patients with non-ST-segment elevation acute coronary syndromes undergoing coronary artery bypass surgery. *J Am Coll Cardiol* 2006;48:281-6.
- Boersma E, Harrington RA, Moliterno DJ, *et al*. Platelet glycoprotein IIb/IIIa inhibitors in acute coronary syndromes: a meta-analysis of all major randomised clinical trials. *Lancet* 2002;359:189-98.
- Pieper KS, Tsiatis AA, Davidian M, *et al*. Differential treatment benefit of platelet glycoprotein IIb/IIIa inhibition with percutaneous coronary intervention versus medical therapy for acute coronary syndromes. *Circulation* 2004;109:641-6.
- Guigliano RP, Newby LK, Harrington RA, *et al*. The early glycoprotein IIb/IIIa inhibition in non-ST-segment elevation acute coronary syndrome (EARLY ACS) trial: a randomized placebo-controlled trial evaluating the clinical benefits of early front-loaded eptifibatide in the treatment of patients with non-ST-segment elevation acute coronary syndromes—study design and rationale. *Am Heart J* 2005;149:994-1002.